

Thrombolysis and Beyond: The New Therapeutic Horizons for Acute Ischemic Stroke

E. Bradshaw Bunney, MD



E. Bradshaw Bunney, MD

Associate Professor
Department of Emergency Medicine
University of Illinois at Chicago
Our Lady of the Resurrection Medical Center
Chicago, IL

E. Bradshaw Bunney, MD



Global Objectives

- Discuss the latest literature and controversy in the use of thrombolysis in stroke
- Discuss options beyond the 3 hour window
- Discuss future therapeutic modalities being studied for the treatment of ischemic stroke

E. Bradshaw Bunney, MD



Clinical History

- 911 call taken by CFD at 2:25 pm
- "My husband is having a stroke and he can not move the left side of his body".
- ALS ambulance arrived at 2:34 pm
- 67-year-old patient to be sitting in a chair with a BP 140/85, pulse 96, respiratory rate 16 and the inability to move his left arm or leg
- His wife also noticed the left side of his face was "flat". He was able to speak.

E. Bradshaw Bunney, MD



Clinical History

- He had a history of hypertension, was on Labetalol and Lasix, with no allergies
- The paramedics noted the time of onset for the symptoms to be 2:15 pm., which was agreed to by both the patient and his wife
- The patient was placed on a cart, an IV was established, oxygen was applied, and glucose was 98
- The patient arrived in the ED at 2:52 pm

E. Bradshaw Bunney, MD




IV Thrombolysis

- The purpose of thrombolysis or clot retrieval in the setting of ischemic stroke is to dissolve or remove the clot
- To preserve the ischemic penumbra and minimize the size of tissue infarct.

E. Bradshaw Bunney, MD





IV Thrombolysis

- By minimizing infarct size
- The NIHSS deficit measured acutely (and long-term in clinical trials)
- The MRS and BI disabilities measured long term can be minimized, improving patient outcomes

E. Bradshaw Bunney, MD

NINDS Trial Results

Percentage with favorable outcome

	Placebo	tPA
No. of patients:	312	145
Modified Rankin Scale	40%	28%
Glasgow Outcome Scale	43%	32%
NIHSS	34%	20%
Symptomatic ICH (within 36 hr)	6.4%	0.6%
Death (by 90 days)	17%	21%

E. Bradshaw Bunney, MD

IV Thrombolysis

- 14% absolute increase for the best clinical outcomes as measured by an NIHSS of 0-1.
- **Benefit** = Need to treat eight patients with tPA in order to have one additional patient with this best outcome.
- 6% absolute increase in the number of symptomatic ICH.
- **Harm** = Will have one symp ICH for every 16 patients treated with tPA.
- 2 patients will have a minimal or no deficit for everyone patient with a symp ICH

E. Bradshaw Bunney, MD

IV Thrombolysis

- In general, tPA should be considered to be optimally useful by reproducing the NINDS protocol
- Studied tPA in patients with a median NIHSS score of 14, signifying a moderate stroke.

E. Bradshaw Bunney, MD

Meta-analyses

E. Bradshaw Bunney, MD

Tale of Meta-analyses

- Wardlaw et al.
- Net benefit despite hazards
- For 1000 treated up to 6hrs, 55 improve, 20 die
- Heterogeneity and wide CI make results unreliable
- Additional trial data required

E. Bradshaw Bunney, MD



Tale of Meta-analyses

- Graham et al., 15 published reports
- ICH rate 5.2%, total death rate 13.4%
- All better than NINDS
- Lysis can be used safely across wide variety of practice settings

E. Bradshaw Bunney, MD



Tale of Meta-analyses

- Hacke et al.
- 6 randomized trials
- Sooner thrombolytics given the greater the benefit
- Particularly when given within 90 min. of onset

E. Bradshaw Bunney, MD



CONTROVERSY: Meta-analysis

- Hoffman and Cooper
- Pooled data can not replace new or confirmatory data
- Meta-analyses did not include streptokinase trials which were negative
- No reason to exclude streptokinase

E. Bradshaw Bunney, MD



Re-analysis

E. Bradshaw Bunney, MD



NINDS Re-analysis

- Does the protocol work?
- Do subgroup imbalances invalidate the entire trial?
- What about BP?


E. Bradshaw Bunney, MD






Baseline NIHSS Imbalance


NIHSS Score		0-5	6-10	11-15	16-20	> 20
No. of patients	Placebo (n=312)	16	83	66	70	77
	t-Pa (n=310)	42	67	65	73	63

Chi-square (4 DF) = 14.8; p = 0.005

E. Bradshaw Bunney, MD 



- ### OTT Analysis Report
- Review Committee had concerns about analyzing OTT as a continuous variable
 - Uncertainty about the exact time of stroke onset.
 - OTT distribution was nonlinear with 25% of all the patients having OTT values of either 89 or 90 minutes.
- E. Bradshaw Bunney, MD 




NINDS ICH Analysis

Risk Factors for ICH:

- Baseline NIHSS > 20
- Age > 70 years
- Ischemic changes present on initial CT
- Glucose > 300 mg/dl (16.7 mmol/L)

# of Risk Factors	# of patients treated with t-PA (n=310)	# Symptomatic ICHs (# of placebo patients with ICH)	Percentage (%)
0	114	2 (1)	1.8
1	144	7 (1)	4.9
> 1	52	11	21.2

E. Bradshaw Bunney, MD 

IV Thrombolysis

- The independent reanalysis of the NINDS tPA clinical trial confirms the results from the initial *NEJM* publication
- Support the use of tPA in stroke patients within three hours of symptom onset
- Number needed to treat calculation based on this reanalysis confirms that approximately 8-10 patients need to be treated with tPA in order to cause one extra patient to have the best clinical outcome.
- 2 patients will improve for every one that develops a symp ICH

E. Bradshaw Bunney, MD



IV Thrombolysis: Conclusion

- tPA has never been demonstrated to be superior or inferior in patients with an acute NIHSS score of 0-5 (mild stroke) or greater than 20 (severe stroke)
- These stroke patients require a more careful assessment of the risks and benefits of tPA
- Since they are less like the patients most commonly treated in the NINDS clinical trial.

E. Bradshaw Bunney, MD



Intra-Arterial Thrombolysis

E. Bradshaw Bunney, MD



IA THROMBOLYSIS

- Two randomized trials – PROACT 1 & 2
- Tested prourokinase vs. heparin <6 hours
- MCA occlusions only
- Recanalization improved with IA
- Mortality identical
- Relative risk reduction for outcome – 60%

E. Bradshaw Bunney, MD



IA Clinical Practice

- Numerous clinical series published
- Basilar artery thrombosis series suggest benefit
- Benefit with basilar may be late (12-24 hrs)
- MRI diffusion/perfusion may aid selection

E. Bradshaw Bunney, MD



IA Thrombolysis

- Within three hours of symptom onset IV tPA is the thrombolytic therapy of choice
- Between three and six hours, there may be a role for intra-arterial tPA in institutions that provide this therapy
- Especially when the stroke is related to occlusion of the middle cerebral artery.

E. Bradshaw Bunney, MD



IA Thrombolysis

- After six hours from stroke symptom onset
- Data suggests that posterior circulation strokes may benefit from attempts to provide intra-arterial thrombolytic therapy
- Data is limited in its scope.

E. Bradshaw Bunney, MD



Future Therapies


E. Bradshaw Bunney, MD



Future Therapies: Neuroprotectants

- First generation failure
- Adverse events
- Lack of efficacy

E. Bradshaw Bunney, MD



E. Bradshaw Bunney, MD



NXY – 059: Preclinical

- Traps carbon and oxygen radicals
- Positive trials in animals/primates
- Significant dose response
- Effective after 4 hours of ischemia

E. Bradshaw Bunney, MD



SAINT I Trial

- Placebo controlled trial
- Acute stroke < 6 hours
- 72 hours infusion of NXY-059
- Primary outcome
 - Disability as measured by Modified Rankin
- BENEFIT at 90 days
 - 4.4% increase in rate of no disability
- No significant AE's

E. Bradshaw Bunney, MD



Future Therapies: Neuroprotectants

- NMDA receptor
- New subtypes
- Antagonists in preclinical trials

E. Bradshaw Bunney, MD



Future Therapies: Neuroprotectants

- Serotonin agonists
- Reduce glutamate-induced excitotoxicity
- Repinotan has reduced infarct volume in preclinical trials
- Up to 5 hours after injury
- Early clinical trials = safe
- Serotonin adverse effects = nausea/vomiting

E. Bradshaw Bunney, MD



Future Therapies: Neuroprotectants

- Inflammatory response in microvasculature
- Leukocyte activation/adhesion
- Good preclinical data, no clinical data of efficacy

E. Bradshaw Bunney, MD



Future Therapies: Hypothermia

- Useful adjunct to other therapies
- Known to be neuroprotective for years
- Positive results in 2 studies with global ischemia
- Timing, degree and duration need further study
- Inconvenient to use

E. Bradshaw Bunney, MD



COOL AID

- 18 patients received hypothermia
- Clinical outcomes similar
- MRI outcomes similar
- Appeared to be well tolerated
- Further studies

E. Bradshaw Bunney, MD



Neuroprotectants

- Neuroprotectants are designed to minimize neuronal cell death and limit infarct size through penumbra stabilization
- A recent *NEJM* publication demonstrated benefit in stroke patients with the use of a novel neuroprotectant
- If confirmed in an ongoing second complementary clinical trial, this would represent the first clinically effective neuroprotectant.

E. Bradshaw Bunney, MD



Informed Consent: Documentation

- With tPA, there is a 30% greater chance of a good outcome at 3 months
- With tPA use, there is 10x greater risk of a symptomatic ICH (severe bleeding stroke)
- Mortality rates at 3 months are the same regardless of whether tPA is used
- 2 patients will have a minimal or no deficit for everyone patient with a symp ICH

E. Bradshaw Bunney, MD



Informed Consent: Documentation

- Patient was explained risks and benefits of tPA use and was able to understand and provide verbal consent (as able), and signature with L hand.
- Risk/benefit favored tPA given clear onset time, young patient with no significant morbidities or factors that would preclude tPA use, and approx NIHSS that suggests OK use.
- Rapid CT obtained, neurology aware of pt status, agreed with expedited tPA use, to follow.

E. Bradshaw Bunney, MD



Documentation

- Just as important
- “The patient is NOT a candidate for tPA because...”

E. Bradshaw Bunney, MD



Case Conclusion

E. Bradshaw Bunney, MD



Clinical Course

- The patient was met by a nurse, a doctor and an EM tech and taken to the resuscitation room.
- They confirmed the onset time of 2:15pm.
- BP 142/88, P 98, R 16, T 99.2 F. HEENT: EOMI, PERRL, Ears clear, neck supple. Heart, lungs and abdomen were normal.
- Neurological exam: CN mild left facial droop, strength 5/5 R arm and leg, 1/5 L arm and leg, no light touch or pin prick sensation in the L arm and leg. NIHSS=17-18.

E. Bradshaw Bunney, MD



Clinical Course

- The stroke team was called at 3:05pm
- Labs were drawn and sent.
- The patient went to CT at 3:20 pm and returned at 3: 41pm.
- The stroke team assessed the patient on return from CT and agreed with the diagnosis of CVA and NIHSS=18.
- Head CT reading was “negative for bleed, normal brain” at 4:03pm.

E. Bradshaw Bunney, MD



Clinical Course

- The patient was felt to be a good candidate for thrombolytics.
- The patient was advised of the risks and benefits.
- The patient, along with his wife declined thrombolytic therapy, stating "I want nature to take its course".
- The patient was given 325 mg. of aspirin and admitted to the hospital.
- His 24 hour NIHSS=14.
- On discharge, 5 days later, NIHSS=10.

E. Bradshaw Bunney, MD



Key Learning Points

- IV thrombolysis is best when used per the NINDS protocol and in patients similar to the NINDS trial
- IA thrombolysis may allow the window to extend to 6 hours in patients with MCA occlusions or posterior circ stroke
- Neuroprotectants may be proven beneficial in the treatment of patients with ischemic stroke in the near future
- Allow patients to make informed decisions

E. Bradshaw Bunney, MD



Questions?

www.ferne.org
ferne@ferne.org

Ferne_2006_AAEM_bunney_thrombolysisFinal

E. Bradshaw Bunney, MD

